Double sections and dominating maps

Gregery T. Buzzard[*] and Steven S. Y. Lu[†]

1 Introduction

As is well-known, given the complex sphere $\mathbb{P}^1$ minus two points, there exist nonconstant holomorphic maps from the plane into this set, the simplest example of which is given by applying the exponential map and then composing with a Mobius transformation taking 0 and ∞ to the two given punctures. Likewise, given the sphere minus one point, we can map the plane into this set by simply applying directly a Mobius transformation taking ∞ to this puncture.

In this paper we prove a parametrized version of this result. To be more precise, we make the following definition.

DEFINITION 1.1 *Let U be a noncompact Riemann surface and let $D \subset U \times \mathbb{P}^1$ be a 1-dimensional complex variety of the form $D = \{(z,w) : a(z)w^2 + b(z)w + c(z) = 0\}$ for holomorphic a, b, c such that for generic $z \in U$ the fiber $D_z = D \cap (\{z\} \times \mathbb{P}^1)$ has two points. Then D is called a double section over U.*

The parametrized version of the above result is that, given a double section D over U, there is a fiber preserving holomorphic map from $U \times \mathbb{C}$ to the complement of D which on each fiber has the form described above.

THEOREM 1.2 *Let D be a double section over U. Then there exists a holomorphic map $F : U \times \mathbb{C} \to (U \times \mathbb{P}^1) \setminus D$ of the form $F(z,w) = (z, H(z,w))$. Moreover, if $\#D_z = 2$, then $H(z,w) = M_z(\exp(c_z w))$, where M_z is a Mobius transformation mapping 0 and ∞ to the two points in D_z and c_z is a nonzero constant depending on z, while if $\#D_z = 1$, then $H(z,w) = M_z(w)$ where M_z is a Mobius transformation mapping ∞ to the point in D_z.*

We will obtain this theorem by first proving a weaker result.

THEOREM 1.3 *Let D be a double section over U. Then there exists a holomorphic function $\sigma : U \to \mathbb{P}^1$ whose graph, a holomorphic section over U, is disjoint from D.*

As a corollary of theorem 1.3 we have the following.

---

[*] Partially supported by an NSF grant.

[†] Partially supported by MSRI and NSERC.

COROLLARY 1.4 *Let D be a double section over U. Then after a biholomorphic change of variables on $U \times \mathbb{P}^1$, there is a holomorphic function h on U such that D has the form*

$$D = \{(z, w) : w^2 = h(z)\}.$$

Although it is possible to obtain these results more abstractly, the methods presented here are completely concrete and allow us to write the maps explicitly in terms of the initial coordinates.

This paper developed out of ideas contained in [BL]. In that paper we showed that for many algebraic surfaces, the existence of a Zariski dense image of $\mathbb{C}$ is equivalent to the existence of a dominating map from $\mathbb{C}^2$ into the surface (a dominating map is a holomorphic map with full rank generically). However, in that paper we did not resolve completely the question of the existence of a Zariski dense image of $\mathbb{C}$ in the complement of a general double section. We resolve the problem completely here.

2 A holomorphic section

In this section we prove theorem 1.3. Let D be a double section in $U \times \mathbb{P}^1$. Since there are only countably many $z \in U$ for which $\#D_z = 1$, we can apply a map of the form $(z, w) \mapsto (z, M(w))$ for some Mobius transformation M to ensure that if $\#D_z = 1$, then $\infty \notin D_z$.

Let E_∞ be the set of $z \in U$ such that $\infty \in D_z$. Then E_∞ is discrete, so we may apply a map of the form $(z, w) \mapsto (z, w - f_0(z))$, where f_0 is holomorphic on U such that if $z \in E_\infty$ and $w \in D_z$, then $w - f_0(z) \neq 0$. Hence if $z_0 \in E_\infty$, there is a neighborhood of z_0 such that in this neighborhood, D is the union of the graphs of two functions of the form $u_0(z)$ and $u_1(z)/(z - z_0)^m$, where $u_j(z_0) \neq 0$ and $m \geq 1$.

Next, let f_1 be holomorphic on U such that if $z_0 \in E_\infty$ with u_0, u_1 as described above, then $f_1(z) = (z - z_0)^m + O(|z - z_0|^{m+1})$. Let $F_1(z, w) = (z, f_1(z)w)$, and let $C = F_1(D)$. Then C is a double section and is contained in $U \times \mathbb{C}$. Moreover, if $z_0 \in E_\infty$, then C is locally the union of the graphs of $v_0(z) = u_0(z)(z - z_0)^m$ and $v_1(z) = u_1(z)$. Moreover, suppose that τ is holomorphic on U such that $\tau(z) \cap C_z = \emptyset$ if $z \notin E_\infty$, while if $z_0 \in E_\infty$, then $\tau(z) = O(|z - z_0|^{m+1})$. Then $F_1^{-1}(z, \tau(z)) = (z, \tau(z)/f_1(z))$ is a holomorphic section which avoids D when $z \notin E_\infty$ and which equals $0 \in \mathbb{C} \setminus \{u_0(z)\}$ when $z \in E_\infty$. Thus $\sigma(z) = \tau(z)/f_1(z)$ gives a holomorphic section in the complement of D, so it suffices to construct τ.

As the final simplification to D, we note that the sum of the points $v_0(z)$ and $v_1(z)$ in C_z is holomorphic on U, so we define the biholomorphic map $F_2(z, w) = (z, w - (v_0(z) + v_1(z))/2)$ and let $B = F_2(C)$. Then B has the form $B = \{(z, w) : w^2 = h(z)\}$, where $h(z) = ((v_0(z) - v_1(z))/2)^2$. Note that if $z_0 \in E_\infty$, then

$$F_2(z, \tau(z)) = \left(z, -\frac{u_0(z)(z - z_0)^m + u_1(z)}{2} + O(|z - z_0|^{m+1}\right), \tag{2.1}$$

so it suffices to construct a section, μ, avoiding B except at points $z_0 \in E_\infty$, at which points μ should have a development as in the second coordinate of (2.1).

To obtain the section μ, we will first define a function $g(z, \sqrt{h(z)})$ which is holomorphic on B except possibly where $h = 0$, then define

$$\mu(z) = \sqrt{h(z)}\frac{g(z, \sqrt{h(z)}) + g(z, -\sqrt{h(z)})}{g(z, \sqrt{h(z)}) - g(z, -\sqrt{h(z)})}. \tag{2.2}$$

Note that for each z, μ is independent of the choice of $\pm\sqrt{h(z)}$, so μ is well-defined and meromorphic except possibly where $h = 0$ or both $g(z, \sqrt{h(z)})$ and $g(z, -\sqrt{h(z)})$ are zero. We will construct g to ensure that at such points, μ has a removable singularity and so that μ avoids B except at points in E_∞, where μ has the form dictated by (2.1).

To this end, let $E_0 = \{z : \#D_z = 1\} = \{z : 0 \in B_z\} = \{z : h(z) = 0\}$. We require

(a) If $z_0 \in E_0$, then $g(z, \sqrt{h(z)}) = 1 + \sqrt{h(z)} + O(\sqrt{|zh(z)|})$.

(b) If $z_0 \in E_\infty$, then taking $\sqrt{h(z)} = (v_0(z) - v_1(z))/2$, we have $g(z, \sqrt{h(z)}) = 1 + z^m + O(|z - z_0|^{m+1})$ and $g(z, -\sqrt{h(z)}) = az^m + O(|z - z_0|^{m+1})$, where $a = u_0(z_0)/u_1(z_0)$.

(c) If $z_0 \notin (E_0 \cup E_\infty)$, then $g(z, \pm\sqrt{h(z)}) \neq 0$.

To obtain such a function g, let $\Psi : X \to B$ be the normalization of B, where X is a noncompact Riemann surface (see e.g. [C]). By [F, theorems 26.3, 26.5], the classical theorems of Weierstrass and Mittag-Leffler are valid on X. Hence given finite holomorphic jets at a discrete set of points in X, there is a function holomorphic on X agreeing with the given jets and having no other zeros. This can be done for instance by first finding a Weierstrass function, f, having the prescribed zeros, dividing all of the jet data by the jets of f at the appropriate points, then taking log of these new jets and interpolating to get a function g having these log jets. Then $f\exp(g)$ has the desired jets and no other zeros. Since Ψ is biholomorphic except where $h = 0$, the data in parts (b) and (c) transfer directly via Ψ to data on X. Hence we need focus only on part (a).

Suppose $z_0 \in E_0$. Without loss we may take $z_0 = 0$. Suppose that $h(z) = z^m h_1(z)$ with m odd and $h_1(0) \neq 0$. Then after a local change of coordinates in X and a choice of $\sqrt{h_1(0)}$, Ψ has the form $\Psi(x) = (x^2, x^m\sqrt{h_1(x^2)})$; i.e., $x = \sqrt{z}$. In this coordinate system, $\Psi^{-1}(z, \pm\sqrt{h(z)}) = \pm x = \pm\sqrt{z}$, and $\sqrt{h(z)} = x^m\sqrt{h_1(x^2)}$. Hence in X, if g_1 is holomorphic in a neighborhood of 0 and satisfies $g_1(x) = 1 + x^m\sqrt{h_1(0)} + O(|x|^{m+1})$, then $g = g_1 \circ \Psi^{-1}$ has the asymptotic development given in (a) in a deleted neighborhood of 0.

Suppose next that $h(z) = z^{2m}h_1(z)$ with $h(0) \neq 0$. In this case, B consists of two distinct components which intersect at the origin, and hence the origin lifts to two distinct points x_0, x_1 in X. Making a choice of $\sqrt{h_1(0)}$, we see that after a local change of coordinates near x_j, Ψ has the form

$$\Psi(x) = (x - x_j, (-1)^j(x - x_j)^m\sqrt{h_1(x - x_j)}).$$

Hence $\Psi^{-1}(z, (-1)^j z^m\sqrt{h_1(z)}) = z + x_j$. As before, in X, if g_1 is holomorphic in a neighborhood of x_j and satisfies $g_1(x) = 1 + (-1)^j(x - x_j)^m\sqrt{h_1(0)} + O(|x - x_j|^{m+1})$, then $g = g_1 \circ \Psi^{-1}$ satisfies the development in (a).

Hence the requirement in part (a) can be satisfied by specifying a discrete set of finite jets in X. As mentioned above, there exists a function g_1 holomorphic on X agreeing with the given finite jets and having no other zeros. Then $g = g_1 \circ \Psi^{-1}$ satisfies (b) and (c) and satisfies (a) in a deleted neighborhood of each $z_0 \in E_0$.

Given g, note that $\mu(z)$ defined as in (2.2) is holomorphic except possibly when $z \in E_0$. Moreover, if z is not in $E_0 \cup E_\infty$, then $g(z, \pm\sqrt{h(z)}) \neq 0$ for either choice of $\pm$, hence $\mu(z) \neq \pm\sqrt{h(z)}$. Hence to finish the proof we need show only that μ is holomorphic and nonzero at each $z_0 \in E_0$, and that μ has the development given by (2.1) at points $z_0 \in E_\infty$.

Suppose first that $z_0 \in E_0$. As before, we may assume $z_0 = 0$ and that $h(z) = z^m h_1(z)$ with $h_1(0) \neq 0$ and $m \geq 1$. Recalling that μ is independent of the choice of $\pm\sqrt{h(z)}$, we have

$$\begin{aligned} \mu(z) &= \sqrt{h}\frac{(1+\sqrt{h}) + (1-\sqrt{h}) + O(|z|^{(m+1)/2})}{(1+\sqrt{h}) - (1-\sqrt{h}) + O(|z|^{(m+1)/2})} \\ &= 1 + O(|z|^{1/2}). \end{aligned}$$

Hence μ has a removable singularity at $z_0 = 0$ and is nonzero there.

Finally, suppose that $z_0 \in E_\infty$, and recall that in this case, $h(z) \neq 0$, and $\sqrt{h(z)} = (v_0(z) - v_1(z))/2$, where, assuming that $z_0 = 0$, we have $v_0(z) = z^m u_0(z)$ and $v_1(z) = u_1(z)$, $u_j(0) \neq 0$, and the development in part (b) holds. Then

$$\begin{aligned} \mu(z) &= \left(\frac{v_0 - v_1}{2}\right)\frac{1 + (1+a)z^m + O(|z|^{m+1})}{1 + (1-a)z^m + O(|z|^{m+1})} \\ &= \left(\frac{u_0 z^m - u_1}{2}\right)(1 + 2az^m + O(|z|^{m+1})) \\ &= -\frac{u_0 z^m + u_1}{2} + u_0 z^m - au_1 z^m + O(|z|^{m+1}). \end{aligned}$$

Since $a = u_0(z_0)/u_1(z_0)$, we see that this last expression has the same form as the second coordinate in (2.1). Thus μ is a holomorphic section with all the desired properties, and hence applying F_2^{-1} and F_1^{-1} gives a holomorphic section avoiding D, as claimed. ■

3 Normal form and dominating map

In this section we prove corollary 1.4 and use this to prove theorem 1.2. The proof of corollary 1.4 is simply a matter of changing coordinates in $U \times \mathbb{P}^1$ so that the section provided by theorem 1.3 becomes the infinity section in the new coordinates. Although this is a standard technique, we provide the proof for completeness.

Proof of corollary 1.4: Let D be a double section over U. We first apply a map which is biholomorphic on $U \times \mathbb{P}^1$ and which maps D into $U \times \mathbb{C}$. To construct such a map, note that by theorem 1.3, there exists a function σ meromorphic on U such that the graph, Σ, of σ in $U \times \mathbb{P}^1$ is disjoint from D. If σ is holomorphic, we apply the map $M(z, w) = (z, 1/(w - \sigma(z)))$, which is biholomorphic on $U \times \mathbb{P}^1$ and which takes Σ to the infinity section and hence satisfies $M(D) \subset U \times \mathbb{C}$.

Otherwise, σ is meromorphic but not holomorphic. Choose a function ψ holomorphic on U such that the zeros of ψ are exactly the zeros of $1/\sigma$, with the same order at each zero. Let $\phi(z) = (-1/\sigma(z)) + (1/g(z))$, and let

$$M(z,w) = \frac{\phi(z)w+1}{w-\sigma(z)}.$$

If $\sigma(z_0) \in \mathbb{C}$, then $\phi(z_0) \in \mathbb{C}$ with $\phi(z_0) \neq -1/\sigma(z_0)$, and hence $M(z_0,\cdot)$ is a nondegenerate Mobius transformation taking $\sigma(z_0)$ to ∞. If σ has a pole of order m at z_0, then ϕ has a pole of the same order at z_0, so multiplying numerator and denominator by $(z-z_0)^m$, we see that $M(z_0,w) = cw$ for some $c \neq 0$. Hence M is biholomorphic and $M(\Sigma)$ is the infinity section, so $M(D)$ is contained in $U \times \mathbb{C}$.

Now, near a generic point in U, $M(D)$ is the union of the graphs of two holomorphic functions u_0 and u_1, and the symmetric function $u_0 + u_1$ is holomorphic on U. Hence the map

$$N(z,w) = \left(z, w - \frac{u_0(z)+u_1(z)}{2}\right)$$

is biholomorphic on $U \times \mathbb{P}^1$, and taking $h(z) = (u_0(z) - u_1(z))^2/4$, we have $N(M(D)) = \{(z,w) : w^2 = h(z)\}$. ■

Finally, we are ready for the construction of a dominating map into the complement of a double section.

Proof of theorem 1.2: Let D be a double section in $U \times \mathbb{P}^1$. By the corollary, we may assume that $D = \{(z,w) : w^2 = h(z)\}$ for some h holomorphic on U. Let

$$F(z,w) = \sqrt{h(z)}\frac{\exp(\sqrt{h(z)}w) + \exp(-\sqrt{h(z)}w)}{\exp(\sqrt{h(z)}w) - \exp(-\sqrt{h(z)}w)}.$$

Note that F is independent of the choice of $\pm\sqrt{h(z)}$. Moreover, F is simply the Mobius transformation $x \mapsto (x+1)/(x-1)$ applied to $\exp(2\sqrt{h(z)}w)$, then multiplied by $\sqrt{h(z)}$. In particular, if $h(z_0) \neq 0$, then $F(z_0,\cdot)$ maps $\mathbb{C}$ into $\mathbb{P}^1 \setminus \{\pm\sqrt{h(z)}\}$, and F is holomorphic except possibly where $h(z) = 0$.

Suppose $h(z_0) = 0$, and without loss take $z_0 = 0$. Then $h(z) = z^m h_1(z)$ for some $m \geq 1$, $h_1(0) \neq 0$. Near 0 we have

$$\begin{aligned} F(z,w) &= \sqrt{h(z)}\frac{(1+\sqrt{h(z)}w) + (1-\sqrt{h(z)}w) + O(|z|^{(m+1)/2})}{(1+\sqrt{h(z)}w) - (1-\sqrt{h(z)}w) + O(|z|^{(m+1)/2})} \\ &= \frac{1+O(|z|^{1/2})}{w + O(|z|^{(m+1)/2})}. \end{aligned}$$

Hence for each fixed w, $F(\cdot,w)$ has a removable singularity at 0, and $F(0,w) = 1/w$. Hence F is holomorphic in each variable separately, hence holomorphic by Hartogs' theorem. Moreover, for each fixed z, F has the form of an exponential map followed by a Mobius transformation if D has two points in the fiber above z, and F has the form of a Mobius transformation if D has one point in the fiber above z. ■

References

[BL] G. Buzzard and S. Lu, Algebraic surfaces holomorphically dominable by $\mathbb{C}^2$, to appear in *Invent. Math.*.

[C] E. Chirka, *Complex analytic sets*, Dordrecht, Boston, Kluwer Academic Publishers, 1989.

[F] O. Forster, *Lectures on riemann surfaces*, Graduate Texts in Mathematics, 81. Springer-Verlag, New York, 1981.

Department of Mathematics
Cornell University
Ithaca, NY 14853 USA

Steven Shin-Yi Lu
Department of Mathematics
University of Waterloo
Waterloo, Ontario, N2L3G1
Canada